

SOME INTERMEDIATE LINEAR ALGEBRA AND BASIC MATRIX THEORY FACTS

DAVID BLECHER
UNIVERSITY OF HOUSTON

MATH 4389

Abstract

This document can be used as a review of some of the things you may not remember from linear algebra (although it does not include some early things that it is assumed that you know, like the meaning of the words in the first few bullets below). Parts of it can be used as a 'formula sheet'. There is some repetition in parts of this document, usually there to help view some things from different angles. There are almost no examples (you do not find examples on a formula sheet), you will have to supply those from the usual sources when you need them. There may be some more simple things that are not said, since we are often focussing here on things that are harder to remember. In places there is a bit more information than you need.

Basics

- It is assumed you know your vectors in \mathbb{R}^n , dot product, their length (Euclidean norm $\|\vec{v}\|_2 = \sqrt{\vec{v} \cdot \vec{v}}$), angle between them, etc. **Cauchy-Schwarz inequality:** $|\vec{a} \cdot \vec{b}| \leq \|\vec{a}\|_2 \|\vec{b}\|_2$.
- You are expected to know the meaning of **size of a matrix** (i.e. $m \times n$) and when two matrices have the same size, or when they are equal, or square ($m = n$).
- It is assumed you know your matrix algebra.
- $A\vec{x} = \sum_k x_k \vec{a}_k$, where \vec{a}_k is the k th column of A , and x_k is the k th entry of \vec{x} . The matrix product

$$AB = [A\vec{b}_1 : A\vec{b}_2 : \cdots : A\vec{b}_r]$$

if \vec{b}_k is the k th column of the $n \times r$ matrix B . Note the i - j entry of AB is the dot product of the i th row of A and the j th column of B .

- You are expected to know the [rules for the transpose](#) A^T , like $(AB)^T = B^T A^T$. We call A symmetric if $A = A^T$.
- You are expected to know a few basic things about [permutation matrices](#) (wiki). These are matrices of 0's and 1's, with exactly one 1 in every column and row.
- You are expected to know what is an upper or lower triangular or diagonal matrix (symbols U, L, D), and what the 'main diagonal' of a matrix is. You may be asked for the LU and maybe the LDU decomposition (wiki), although this is not likely.
- We write I or I_n for the $n \times n$ [identity matrix](#). This is both a diagonal matrix and a permutation matrix. Note $A = AI = IA$.

E.g. $I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$.

- We will discuss the **inverse** A^{-1} of a square matrix A in more detail later, but for now: a matrix is **nonsingular** iff it is invertible (has an inverse, that is, a matrix B with $AB = BA = I$. We write such B , if it exists, as A^{-1}). Otherwise, A is called **singular**. The inverse of a diagonal matrix $\text{diag}(d_1, d_2, \dots, d_n)$ is $\text{diag}(d_1^{-1}, d_2^{-1}, \dots, d_n^{-1})$, assuming the d_k are all nonzero.

- The inverse of a 2×2 matrix:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix},$$

assuming that the determinant $ad - bc \neq 0$. (If $ad - bc = 0$ the matrix is not invertible.)

- For a general 3×3 or larger square matrix A you can find the inverse A^{-1} if it is invertible either by (1) the cofactor method using the determinant (wiki), or (2): Try to use Gauss-Jordan elimination (below) on the matrix $[A : I]$ to achieve reduced row echelon form $[I : C]$. If you can do this then $C = A^{-1}$, otherwise A is not invertible.

- If A is an $m \times n$ matrix, we can use Gauss elimination to row reduce A to row echelon form (REF) (or 'staircase form' see pic). We can also use Gauss-Jordan elimination to row reduce A to reduced row echelon form (RREF). By an echelon form of A we mean either of the above.

We will write $\text{RREF}(A)$ for the reduced row echelon form. Convention: we will make $\text{RREF}(A)$ have 1's in all the pivots (so that each pivot column has one 1 and the rest zeros).

- One may view the row echelon form or the RREF of A as the product EA of a matrix E and A . Indeed E is the product of 'elementary matrices' each representing and performing (by left multiplying by it, in the right order) one of the *elementary row operations* needed in the row reduction.
[Picture drawn of matrix in ('staircase') form REF or RREF]

- There are **three elementary row operations**, namely switching two rows, multiplying one row by a nonzero scalar k , or adding to one row a nonzero multiple of another row.

Hence there are three kinds of elementary matrices (you will have seen them, write them down), all invertible. The first is a simple permutation matrix (which is its own inverse). Indeed the elementary matrix that corresponds to switching rows i and j is the identity matrix with rows i and j switched. The second is a simple diagonal matrix which looks like the identity matrix except for a k in one of the diagonal entries (and its inverse is the same except for a $1/k$ in that entry). The third kind, adding to row j a nonzero multiple k of row i is a lower triangular matrix which looks like the identity matrix except for a k in the $j-i$ entry (and its inverse is the same except for a $-k$ in that spot).

- You are expected to know the meaning of ‘pivot positions’, ‘pivot columns’ and ‘free variables’ (the variables corresponding to the non-pivot columns). Non-pivot columns are sometimes called free columns. Pictures shown in class.
- The augmented matrix of a linear system $A\vec{x} = \vec{b}$ is the matrix $[A : \vec{b}]$.
- We usually view vectors in \mathbb{R}^n as columns (as in the last line). Sometimes to stress this point we call them column vectors.

- We will not define a **vector space** here. If you have not looked at that for a while do so now, eg. on wikipedia, and be sure that you have a feeling for which examples are vector spaces and which are not.

The main vector spaces we look at in this course are \mathbb{R}^n , and its subspaces (particularly the ones met with below). By a **scalar** we mean a real number (although nearly everything here applies to complex vector spaces too).

- Recall that a **subspace** of a vector space V is a subset W of V containing the 0 vector of V , such that (a) $v + w \in W$, and (b) $cv \in W$, for all $v, w \in W$ and for all scalars c .

FACT: Any subspace of a vector space is a vector space.

- A **linear combination** of vectors v_1, v_2, \dots, v_m is a vector of form

$$c_1v_1 + c_2v_2 + \dots + c_mv_m$$

for scalars c_1, \dots, c_m . Here $m \in \mathbb{N}$.

- Writing one vector \vec{b} in \mathbb{R}^n as a linear combination of several other vectors: Number these other vectors, and let A be the matrix having these vectors as columns in this order. Solve $A\vec{x} = \vec{b}$. The coefficients in the desired linear combination are the entries in any particular solution of this equation.
- Recall that the solution of $A\vec{x} = \vec{b}$ is $\vec{x} = A^{-1}\vec{b}$ if A is square and invertible.
- The span of a set B of vectors is the set of all linear combinations of vectors in B ; it is written $\text{Span}(B)$. **FACT:** $\text{Span}(B)$ is a subspace.

We say a set B of vectors in a vector space V spans V , or is a spanning set for V , if $\text{Span}(B) = V$.

- A set $B = \{v_1, v_2, \dots, v_m\}$ in a vector space V is said to be **linearly dependent** if one of them can be written as a linear combination of the others. Otherwise, B is said to be **linearly independent** or **l.i.** for short.

A set of one nonzero vector $\{v\}$ is always linearly independent. Two vectors are linearly dependent if and only if one is a scalar multiple of the other. Two vectors in \mathbb{R}^n are linearly independent if and only if they are not parallel (and neither is $\vec{0}$). Three vectors in \mathbb{R}^3 are linearly independent if and only if they do not lie in the same plane.

- **FACT:** $B = \{v_1, v_2, \dots, v_m\}$ is linearly independent if and only if the only scalars c_1, c_2, \dots, c_m such that $c_1v_1 + c_2v_2 + \dots + c_mv_m = \vec{0}$ are $c_1 = c_2 = \dots = c_m = 0$.

- **Test** if a collection of several vectors in \mathbb{R}^n is linearly independent: Let A be the matrix having these vectors as columns. If this (or its RREF) has free variables then the vectors are linearly dependent.

Saying these vectors are linearly independent is the same as saying that $A\vec{x} = \vec{0}$ has only the trivial solution $\vec{0}$.

- A **basis** for a vector space V is a finite set in V which is both linearly independent and spans V .
- A finite set $B = \{v_1, v_2, \dots, v_m\}$ in V is a basis for V iff every vector v in V can be written as a linear combination $v = c_1v_1 + c_2v_2 + \dots + c_mv_m$ in one and only one way (that is, c_1, \dots, c_m are unique). We call this 'one way' *writing (or expressing) v in terms of the basis*.
- A vector space V is **finite dimensional** if it has a (finite) basis. In this case one can show that all bases of V are the same size. This size is called the **dimension** of V , and is written as $\dim(V)$. If a vector space V is not finite dimensional it is called **infinite dimensional** (and then we will not talk about bases for V).
- **FACT:** The dimension of the span of a set S of several vectors in \mathbb{R}^n is the maximum number of linearly independent members of S . We will give a recipe later.
- **FACT:** A strictly smaller subspace (that is $W \subset V$ but $W \neq V$) will have strictly smaller dimension. (Assuming finite dimensions.)

- Suppose that $\dim(V) = n$. Any set of strictly more than (resp. less than) n vectors in V is linearly dependent (resp. cannot span V). Any linearly independent (resp. spanning) set in V has $\leq n$ (resp. $\geq n$) elements.

So any set of linearly independent vectors in \mathbb{R}^n has $\leq n$ members, any spanning set for \mathbb{R}^n has $\geq n$ members.

- **FACT:** A set of n vectors in \mathbb{R}^n is linearly independent if and only if it spans \mathbb{R}^n . That is, if and only if it is a basis for \mathbb{R}^n . This is also equivalent to saying that the matrix with these vectors as columns is invertible (or that this matrix has nonzero determinant).
- The **standard basis** or **canonical basis** for \mathbb{R}^3 from Calculus III is often written as $\vec{i}, \vec{j}, \vec{k}$. The **standard basis** or **canonical basis** for \mathbb{R}^n is often written as $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$.

- Column space $C(A)$ of a matrix A :
- If A is an $m \times n$ matrix, we can write $A = [\vec{a}_1 : \vec{a}_2 : \cdots : \vec{a}_n]$; here \vec{a}_k is the k th column of A . Then $C(A)$ is the set of all the linear combinations of $\vec{a}_1, \vec{a}_2, \cdots, \vec{a}_n$. So we really have $C(A) = \text{Span}(\vec{a}_1, \vec{a}_2, \cdots, \vec{a}_n)$. This is a vector space, is a subspace of \mathbb{R}^m as we said above.
- Also, $C(A) = \{\vec{b} \in \mathbb{R}^m : A\vec{x} = \vec{b} \text{ has a solution}\}$. That is, asking if $\vec{b} \in C(A)$ is the same as asking: does $A\vec{x} = \vec{b}$ have a solution?
 (To prove this, recall that for any $\vec{x} \in \mathbb{R}^n$, we can write $A\vec{x} = x_1 \vec{a}_1 + x_2 \vec{a}_2 + \cdots + x_n \vec{a}_n$ (here x_k is the k th entry of \vec{x}). Thus saying that $A\vec{x} = \vec{b}$ has a solution, is the same as saying that \vec{b} is a linear combination of the columns of A , or equivalently, as saying that $\vec{b} \in \text{Span}(\vec{a}_1, \vec{a}_2, \cdots, \vec{a}_n)$.)
- So one way to find $C(A)$ is to use Gauss elimination to find the \vec{b} such that $A\vec{x} = \vec{b}$ has a solution.

- Finding a **nice basis** for $C(A)$: Do Gauss-Jordan to A^T to obtain $\text{RREF}(A^T)$. Take the transpose of $\text{RREF}(A^T)$. The nonzero columns of this, that is the nonzero columns of $(\text{RREF}(A^T))^T$, form a basis for $C(A)$.
- Finding a basis for $C(A)$ **consisting of columns** of A : Do Gauss elimination to A . The original columns of A standing in the pivot columns, are the desired basis.
- **Finding a basis for the span of several vectors** in \mathbb{R}^n : Number them. Let A be the matrix having these vectors as columns in this order, and apply one of the last two items to find a basis for $C(A)$.

- Given a finite set of vectors S find a subset which is a basis for $\text{Span}(S)$: this is the same recipe as for the last item, but using the item before it to find a basis for $C(A)$.
- **Row space** $R(A)$ of a matrix A is the span of the rows of A . Note that $R(A) = C(A^T)$ and $C(A) = R(A^T)$.
- **Finding a basis for $R(A)$** : Do Gauss-Jordan to A . The nonzero rows in $\text{RREF}(A)$ are a basis for $R(A)$. Or you could find a basis for $C(A^T)$ by one of the methods above.
- $\text{rank}(A) = \text{number of nonzero rows in } \text{RREF}(A) = \text{number of pivot columns/pivot variables} = \dim(C(A)) = \dim(R(A))$.
- **FACT**: $\text{rank}(A) = \text{rank}(A^T)$.
- A **rank 1 matrix** may also be described as a matrix of the form $A = \vec{v} \vec{w}^T$, for vectors \vec{v}, \vec{w} .

- The **nullspace** $N(A)$ for matrix A is the set of solutions to the homogeneous equation $A\vec{x} = \vec{0}$. It is a subspace, and always contains the zero vector $\vec{0}$, the trivial solution to $A\vec{x} = \vec{0}$. The nullspace is also the **kernel** $\text{Ker}(L_A)$ of the operator $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ of left multiplication by A on (column) vectors in \mathbb{R}^n . Then **nullity** $(A) = \dim N(A)$.
- **FACT:** If A has more columns than rows then $A\vec{x} = \vec{0}$ has infinitely many solutions, and $\text{nullity}(A) \geq 1$.
- **Rank-nullity theorem:** $\text{rank}(A) + \text{nullity}(A) = n$ for an $m \times n$ matrix A .
(Sketch proof: this is almost obvious for $\text{RREF}(A)$, and none of the operations for RREF change rank, nullity, n .)

- The members of a basis for $N(A)$ are called the **fundamental solutions**.
- The **special solutions** are a particularly ‘nice’ basis for $N(A)$. Here is the recipe for finding the special solutions. First compute $\text{RREF}(A)$. Then each special solution is found by solving $\text{RREF}(A) \vec{x} = \vec{0}$, putting all the free variables equal to zero except one, which we set equal to 1. (It is ‘nice’ because of all the ‘1’s in positions where all other vectors have 0’s.)

The **number of fundamental solutions** is the number of free variables in an echelon form of A . This number is the **dimension of $N(A)$** .

- If $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_m$ are the fundamental solutions (i.e. any basis for $N(A)$) then the **general solution** to $A\vec{x} = \vec{0}$ is

$$c_1\vec{x}_1 + c_2\vec{x}_2 + \dots + c_m\vec{x}_m$$

for scalars c_1, c_2, \dots, c_m .

(This follows from the definition of a basis.)

- The general solution to $A\vec{x} = \vec{b}$ is

$$\vec{x} = \vec{x}_h + \vec{x}_p = \vec{x}_p + c_1\vec{x}_1 + c_2\vec{x}_2 + \cdots + c_m\vec{x}_m.$$

Here \vec{x}_h is the general solution to the associated homogeneous equation $A\vec{x} = \vec{0}$, and \vec{x}_p is a particular solution to $A\vec{x} = \vec{b}$, and $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_m$ are the fundamental solutions to the associated homogeneous equation $A\vec{x} = \vec{0}$ (that is, they are a basis for $N(A)$).

One way to find a particular solution to $A\vec{x} = \vec{b}$ is to do Gauss-Jordan elimination to the augmented matrix, then set the free variables equal to 0.

Theorem The general solution to $A\vec{x} = \vec{b}$ is $\vec{x} = \vec{x}_h + \vec{x}_p$ where \vec{x}_h is the general solution to the associated homogeneous equation $A\vec{x} = \vec{0}$, and \vec{x}_p is a particular solution to $A\vec{x} = \vec{b}$.

Proof: I). $\vec{x}_h + \vec{x}_p$ is a solution to $A\vec{x} = \vec{b}$:

$$A(\vec{x}_h + \vec{x}_p) = A\vec{x}_h + A\vec{x}_p = \vec{0} + \vec{b} = \vec{b}.$$

II). Now suppose that \vec{x}_0 is any solution to $A\vec{x} = \vec{b}$. Then

$$A(\vec{x}_0 - \vec{x}_p) = A\vec{x}_0 - A\vec{x}_p = \vec{b} - \vec{b} = \vec{0}.$$

Thus $\vec{x}_0 - \vec{x}_p$ solves the associated homogeneous equation $A\vec{x} = \vec{0}$, so $\vec{x}_0 - \vec{x}_p$ is of the form \vec{x}_h .

But if $\vec{x}_0 - \vec{x}_p = \vec{x}_h$ then $\boxed{\vec{x}_0 = \vec{x}_h + \vec{x}_p}$ □

Key tool above: **LINEARITY** $A\vec{x}$ is a linear transformation of \vec{x} .

Theorem The general solution to $A\vec{x} = \vec{b}$ is

$$\vec{x} = \vec{x}_h + \vec{x}_p = \boxed{\vec{x}_p + c_1\vec{x}_1 + c_2\vec{x}_2 + \cdots + c_m\vec{x}_m}.$$

Here \vec{x}_h, \vec{x}_p are as before, and $\vec{x}_1, \vec{x}_2, \cdots, \vec{x}_m$ are the **fundamental solutions** to the associated homogeneous equation $A\vec{x} = \vec{0}$.

Proof: Combine the last theorem and a fact a few pages back (after we defined the fundamental solutions) that says $\vec{x}_h = c_1\vec{x}_1 + c_2\vec{x}_2 + \cdots + c_m\vec{x}_m$. So by the last theorem the general solution to $A\vec{x} = \vec{b}$ is

$$\vec{x} = \vec{x}_h + \vec{x}_p = \vec{x}_p + c_1\vec{x}_1 + c_2\vec{x}_2 + \cdots + c_m\vec{x}_m. \quad \square$$

Finding solutions to linear equations or systems is close to the heart of linear algebra.

There is a similar fact in differential equations, which has the 'same' proof: Suppose that L is a linear differential operator (like $3D^2 - e^t D + 5I$), that x_h is the general solution to the associated homogeneous equation $Lx = 0$, and that x_p is a particular solution to $Lx = g$. Show that the general solution to $Lx = g$ is $x = x_p + x_h$. And $x_h = c_1x_1 + c_2x_2 + \cdots + c_mx_m$ for scalars c_1, c_2, \cdots, c_m , where x_1, x_2, \cdots, x_m are the fundamental solutions, i.e. a basis for the solution space of the associated homogeneous equation $Lx = 0$.

Proof: That $x_h = c_1x_1 + c_2x_2 + \cdots + c_mx_m$ follows as before from the definition of basis and that x_1, x_2, \cdots, x_m are the fundamental solutions, i.e. a basis for the solution space of the associated homogeneous equation $Lx = 0$.

I). $x_h + x_p$ is a solution to $Lx = g$:

$$L(x_h + x_p) = Lx_h + Lx_p = 0 + g = g.$$

II). Now suppose that x_0 is any solution to $Lx = g$. Then

$$L(x_0 - x_p) = Lx_0 - Lx_p = g - g = 0.$$

Thus $x_0 - x_p$ solves the associated homogeneous equation $Lx = 0$, so $x_0 - x_p$ is of the form x_h .

But if $x_0 - x_p = x_h$ then $x_0 = x_h + x_p$. □

Key tool above: LINEARITY L is a linear transformation .

Table of the number of solutions to $A\vec{x} = \vec{b}$ for an $m \times n$ matrix A :

If $m = n$ there is 1 solution if A is invertible; if A is not invertible then there are zero or infinitely many solutions.

If $m > n = \text{rank}(A)$ there are 1 or no solutions.

If $m < n$ there are none or infinitely many solutions (there are infinitely many if $\text{rank}(A) = m$).

If $\text{rank}(A) < \min\{n, m\}$ there are none or infinitely many solutions. (All of these can be seen/proved by thinking about RREF of the augmented matrix, and/or using the second last bullet).

- On inverses: If $AB = I$ or $BA = I$ and A and B have the same size then $A = B^{-1}$ (and $B = A^{-1}$).
- If A and B are invertible matrices of the same size then the inverse of AB is $B^{-1}A^{-1}$.
- If A is an invertible matrix then A^T has inverse $(A^{-1})^T$.

- **Determinants:** only make sense for square matrices.
- if A is a square matrix then A is **invertible** iff $\det(A) \neq 0$.
- You should know the **Cramer's rule** method for solving the system $A\vec{x} = \vec{b}$ if A is a square matrix.
- You are expected to know the **main properties of determinants**: like $\det(A) = \det(A^T)$, $\det(A^{-1}) = 1/\det(A)$, $\det(AB) = \det(A)\det(B)$, etc.
- Or, that the determinant changes sign if you switch two rows or columns, is multiplied by k if you multiply a row (or column) through by k , and is unchanged if you add to one row a nonzero multiple of another row . If one row (resp. column) of a matrix equals or is a multiple of another row (resp. column) its determinant is zero. Etc.
- You are expected to know the formula for the determinant of a 2×2 matrix ($ad - bc$), or of an upper or lower triangular matrix (the product of the main diagonal entries).

- You are expected to know the 4 methods for the determinant of a 3×3 matrix, three of which work for bigger square matrices too (cofactors, Gauss elimination to lower triangular, the permutation definition).
- You should know the connections between determinants and the cross product of vectors, or the volume of a parallelepiped (google this).

Theorem If A is an $n \times n$ matrix the following are equivalent:

- (1) A is invertible.
- (2) $\det(A) \neq 0$.
- (3) $\text{rank}(A) = n$.
- (4) For every vector \vec{b} the system $A\vec{x} = \vec{b}$ has a solution.
- (5) For every vector \vec{b} the system $A\vec{x} = \vec{b}$ has a unique solution.
- (6) For some vector \vec{b} the system $A\vec{x} = \vec{b}$ has a unique solution.
- (7) The system $A\vec{x} = \vec{0}$ has only the trivial solution $\vec{x} = \vec{0}$ (i.e. $N(A) = \{\vec{0}\}$).
- (8) The columns of A are linearly independent.
- (9) The columns of A span n -space.
- (10) The rows of A are linearly independent.
- (11) The rows of A span n -space.
- (12) The reduced row echelon form of A is I_n .

(13) The (echelon forms) of A have n pivot positions/columns.

(14) There is a matrix B with $AB = I$.

(15) There is a matrix B with $BA = I$.

Diagonalization

An n by n matrix can be diagonalized (that is, $A = SDS^{-1}$ for a diagonal matrix D and an invertible matrix S) if and only there exist n linearly independent eigenvectors, and if and only if the algebraic multiplicity of each eigenvalue equals its geometric multiplicity.

If S is the matrix with these n linearly independent eigenvectors as columns, and if D is a diagonal matrix with the associated (all real) eigenvalues (in the same order as the column eigenvectors) as diagonal entries, then $A = SDS^{-1}$.

Orthogonal diagonalization

An n by n real matrix is symmetric (i.e. $A = A^T$) if and only there exist n mutually orthogonal real eigenvectors, and if and only if there is an orthonormal basis of real eigenvectors of A , and if and only if $A = UDU^T$ for real diagonal matrix D and real orthogonal matrix U .

If U is the matrix with these n orthonormal real eigenvectors as columns, and if D is a diagonal matrix with the associated (all real) eigenvalues (in the same order as the column eigenvectors) as diagonal entries, then $A = UDU^T$.

There is a similar list of equivalent conditions for an $m \times n$ matrix, for when $\text{rank}(A) = m$ etc (and another such list when $\text{rank}(A) = n$).

The rest of the ' Linear algebra 'important facts' list (partial)' pdf document linked on the 'Notes for most of the modules' page, was not covered in class.